

Title: Nephronophthisis GeneReview (Overview) – Molecular Genetics: Animal Models
Authors: Stokman N, Lilien M, Knoers N

Posted: June 2016

Note: The following information is provided by the authors and has not been reviewed by GeneReviews staff.

Animal Models for Genes Associated with Nephronophthisis

[NPHP1](#)

[INVS \(NPHP2\)](#)

[NPHP3](#)

[NPHP4](#)

[NPHP5 \(IQCB1\)](#)

[NPHP6 \(CEP290\)](#)

[TMEM67 \(NPHP11\)](#)

NPHP1

Loss of *NPHP1* in Madin-Darby canine kidney epithelial (MDCK) cells resulted in reduced cilia formation and length, delayed tight-junction formation, and disorganized multi-lumen cyst formation [Delous et al 2009]. A mouse model with homozygous deletions of exon 20 of *NPHP1* showed retinal degeneration with abnormal photoreceptor intraflagellar transport [Jiang et al 2009]. Interestingly, the mice did not have nephronophthisis but did show impaired spermatogenesis [Jiang et al 2008].

INVS (NPHP2)

Mouse embryonic fibroblasts (MEF) deficient for INVS showed normal cilia formation and cilia length but defective cilia orientation and directional cell migration [Veland et al 2013]. A mouse model of *Inv* mutation showed situs inversus and cystic renal disease [Yokoyama et al 1993, Morgan et al 1998]. Morpholino knockdown of Invs in zebrafish resulted in ventral axis curvature, left-right asymmetry and pronephric cysts [Simons et al 2005, Zhao & Malicki 2011].

NPHP3

Knockdown of *NPHP3* in a murine inner medullary collecting duct (IMCD3) cell model resulted in a reduced number of cilia and disrupted spheroid formation caused by polarity defects, effects that could be rescued by the somatostatin agonist octreotide [Ghosh et al 2012].

Homozygous pathogenic variants in *Nphp3* were responsible for the phenotype in the pcy (polycystic kidney disease) mouse mutant [Olbrich et al 2003].

Homozygous null-mutations resulted in lateralization defects and were embryonically lethal in mice [Bergmann et al 2008].

Morpholino knockdown of *Nphp3* in zebrafish embryos caused abnormal body curvature, hydrocephalus, pronephric cysts, and situs inversus [Zhou et al 2010, Hoff et al 2013].

NPHP4

NPHP4 knockdown in Madin-Darby canine kidney (MDCK) cells resulted in significantly shorter cilia, delayed tight-junction formation, and disorganized multi-lumen structures when grown in three-dimensional collagen matrix [Delous et al 2009]. Unlike *NPHP1* knockdown, silencing of *NPHP4* did not lead to a reduced number of cilia [Delous et al 2009].

Morpholino knockdown of *Nphp4* in zebrafish caused abnormal body curvature, laterality defects, hydrocephalus, pericardial edema, retinal abnormalities, pronephric cysts and cloaca defects [Burcklé et al 2011, Slanchev et al 2011, Borgal et al 2012, French et al 2012].

Mouse KO, retinal involvement, and male infertility [Won et al 2011].

NPHP5 (IQCB1)

Depletion of *lqcb1* in zebrafish results in abnormal body curvature, cerebral anomalies, and development of pronephric cysts. This phenotype is aggravated by the combined knockdown of *CEP290*, suggesting a genetic interaction [Schäfer et al 2008].

NPHP6 (CEP290)

Knockdown of *CEP290* in an IMCD3 cell model caused reduced cilia formation and increased abnormal spheroid formation [Ghosh et al 2012].

Knockdown of *Cep290* in zebrafish caused abnormal body curvature, cerebral and retinal anomalies and development of pronephric cysts [Schäfer et al 2008].

Cep290 knockout in mice recapitulated the cerebellar Joubert phenotype and resulted in retinal and renal pathology [Lancaster et al 2011, Hynes et al 2014, Rachel et al 2015].

TMEM67 (NPHP11)

Morpholino knockdown of *Tmem67* in zebrafish resulted in a ciliopathy phenotype characterized by brain and eye abnormalities and pronephric cyst formation. Concomitant knockdown of *Flna* aggravated this phenotype [Adams et al 2012].

Tmem67 knockdown mice manifested renal cystic dysplasia, neurologic features associated with ciliopathies, limb dysplasia, laterality defects, cardiac defects, pulmonary hypoplasia, and defective Wnt and Shh signaling [Abdelhamed et al 2013, Abdelhamed et al 2015].

References

- Abdelhamed ZA, Natarajan S, Wheway G, Inglehearn CF, Toomes C, Johnson CA, Jagger DJ. The Meckel-Gruber syndrome protein TMEM67 controls basal body positioning and epithelial branching morphogenesis in mice via the non-canonical Wnt pathway. *Dis Model Mech.* 2015;8:527-41.
- Abdelhamed ZA, Wheway G, Szymanska K, Natarajan S, Toomes C, Inglehearn C, Johnson CA. Variable expressivity of ciliopathy neurological phenotypes that encompass Meckel-Gruber syndrome and Joubert syndrome is caused by complex de-regulated ciliogenesis, Shh and Wnt signalling defects. *Hum Mol Genet.* 2013;22:1358-72.
- Adams M, Simms RJ, Abdelhamed Z, Dawe HR, Szymanska K, Logan CV, Wheway G, Pitt E, Gull K, Knowles MA, Blair E, Cross SH, Sayer JA, Johnson CA. A meckelin-filamin A interaction mediates ciliogenesis. *Hum Mol Genet.* 2012;21:1272-86.
- Bergmann C, Fliegauf M, Brüchle NO, Frank V, Olbrich H, Kirschner J, Schermer B, Schmedding I, Kispert A, Kränzlin B, Nürnberg G, Becker C, Grimm T, Girschick G, Lynch SA, Kelehan P, Senderek J, Neuhaus TJ, Stallmach T, Zentgraf H, Nürnberg P, Gretz N, Lo C, Lienkamp S, Schäfer T, Walz G, Benzing T, Zerres K, Omran H. Loss of nephrocystin-3 function can cause embryonic lethality, Meckel-Gruber-like syndrome, situs inversus, and renal-hepatic-pancreatic dysplasia. *Am J Hum Genet.* 2008;82:959-70.
- Borgal L, Habbig S, Hatzold J, Liebau MC, Dafinger C, Sacarea I, Hammerschmidt M, Benzing T, Schermer B. The ciliary protein nephrocystin-4 translocates the canonical Wnt regulator Jade-1 to the nucleus to negatively regulate beta-catenin signaling. *J Biol Chem.* 2012;287:25370-80.
- Burcklé C, Gaudé HM, Vesque C, Silbermann F, Salomon R, Jeanpierre C, Antignac C, Saunier S, Schneider-Maunoury S. Control of the Wnt pathways by nephrocystin-4 is required for morphogenesis of the zebrafish pronephros. *Hum Mol Genet.* 2011;20:2611-27.
- Delous M, Hellman NE, Gaudé HM, Silbermann F, Le Bivic A, Salomon R, Antignac C, Saunier S. Nephrocystin-1 and nephrocystin-4 are required for epithelial morphogenesis and associate with PALS1/PATJ and Par6. *Hum Mol Genet.* 2009;18:4711-23.
- French VM, Van de Laar IMBH, Wessels MW, Rohe C, Roos-Hesselink JW, Wang G, Frohn-Mulder IM, Severijnen LA, de Graaf BM, Schot R, Breedveld G, Mientjes E, van Tienhoven M, Jadot E, Jiang Z, Verkerk A, Swagemakers S, Venselaar H, Rahimi Z, Najmabadi H, Meijers-Heijboer H, de Graaff E, Helbing WA, Willemse R, Devriendt K, Belmont JW, Oostra BA, Amack JD, Bertoli-Avella AM. NPHP4 variants are associated with pleiotropic heart malformations. *Circ Res.* 2012;110:1564-74.
- Ghosh AK, Hurd T, Hildebrandt F. 3D spheroid defects in NPHP knockdown cells are rescued by the somatostatin receptor agonist octreotide. *Am J Physiol Renal Physiol.* 2012;303:F1225-9.
- Hoff S, Halbritter J, Epting D, Frank V, Nguyen TM, van Reeuwijk J, Boehlke C, Schell C, Yasunaga T, Helmstädtler M, Mergen M, Filhol E, Boldt K, Horn N, Ueffing M, Otto EA, Eisenberger T, Elting MW, van Wijk JA, Bockenhauer D, Sebire NJ, Rittig S, Vyberg M, Ring T, Pohl M, Pape L, Neuhaus TJ, Elshakhs NA, Koon SJ, Harris PC, Grahammer F, Huber TB, Kuehn EW, Kramer-Zucker A, Bolz HJ, Roepman R, Saunier S, Walz G, Hildebrandt F, Bergmann C, Lienkamp SS. ANKS6 is a central component of a nephronophthisis module linking NEK8 to INVS and NPHP3. *Nat Genet.* 2013;45:951-6.
- Hynes AM, Giles RH, Srivastava S, Eley L, Whitehead J, Danilenko M, Raman S, Slaats GG, Colville JG, Ajzenberg H, Kroes HY, Thelwall PE, Simmons NL, Miles CG, Sayer JA. Murine Joubert syndrome

reveals Hedgehog signaling defects as a potential therapeutic target for nephronophthisis. *Proc Natl Acad Sci USA* 2014;111:9893-8.

Jiang ST, Chiou YY, Wang E, Lin HK, Lee SP, Lu HY, Wang CK, Tang MJ, Li H. Targeted disruption of Nphp1 causes male infertility due to defects in the later steps of sperm morphogenesis in mice. *Hum Mol Genet*. 2008;17:3368-79.

Lancaster MA, Gopal DJ, Kim J, Saleem SN, Silhavy JL, Louie CM, Thacker BE, Williams Y, Zaki MS, Gleeson JG. Defective Wnt-dependent cerebellar midline fusion in a mouse model of Joubert syndrome. *Nat Med*. 2011;17:726-31.

Morgan D, Turnpenny L, Goodship J, Dai W, Majumder K, Matthews L, Gardner A, Schuster G, Vien L, Harrison W, Elder FF, Penman-Splitt M, Overbeek P, Strachan T. Inversin, a novel gene in the vertebrate left-right axis pathway, is partially deleted in the inv mouse. *Nat Genet*. 1998;20:149-56.

Olbrich H, Fliegauf M, Hoefele J, Kispert A, Otto E, Volz A, Wolf MT, Sasmaz G, Trauer U, Reinhardt R, Sudbrak R, Antignac C, Gretz N, Walz G, Schermer B, Benzing T, Hildebrandt F, Omran H. Mutations in a novel gene, NPHP3, cause adolescent nephronophthisis, tapeto-retinal degeneration and hepatic fibrosis. *Nat Genet*. 2003;34:455-9.

Rachel RA, Yamamoto EA, Dewanjee MK, May-Simera HL, Sergeev YV, Hackett AN, Pohida K, Munasinghe J, Gotoh N, Wickstead B, Fariss RN, Dong L, Li T, Swaroop A. CEP290 alleles in mice disrupt tissue-specific cilia biogenesis and recapitulate features of syndromic ciliopathies. *Hum Mol Genet*. 2015;24:3775-91.

Schäfer T, Putz M, Lienkamp S, Ganner A, Bergbreiter A, Ramachandran H, Gieloff V, Gerner M, Mattonet C, Czarnecki PG, Sayer JA, Otto EA, Hildebrandt F, Kramer-Zucker A, Walz G. Genetic and physical interaction between the NPHP5 and NPHP6 gene products. *Hum Mol Genet*. 2008;17:3655-62.

Slanchev K, Putz M, Schmitt A, Kramer-Zucker A, Walz G. Nephrocystin-4 is required for pronephric duct-dependent cloaca formation in zebrafish. *Hum Mol Genet*. 2011;20:3119-28.

Veland IR, Montjean R, Eley L, Pedersen LB, Schwab A, Goodship J, Kristiansen K, Pedersen SF, Saunier S, Christensen ST. Inversin/Nephrocystin-2 is required for fibroblast polarity and directional cell migration. *PLoS One*. 2013;8:e60193.

Won J, Marín de Evsikova C, Smith RS, Hicks WL, Edwards MM, Longo-Guess C, Li T, Naggert JK, Nishina PM. NPHP4 is necessary for normal photoreceptor ribbon synapse maintenance and outer segment formation, and for sperm development. *Hum Mol Genet*. 2011;20:482-96.

Yokoyama T, Copeland NG, Jenkins NA, Montgomery CA, Elder FF, Overbeek PA. Reversal of left-right asymmetry: a situs inversus mutation. *Science*. 1993;260:679-82.

Zhao C, Malicki J. Nephrocystins and MKS proteins interact with IFT particle and facilitate transport of selected ciliary cargos. *EMBO J*. 2011;30:2532-44.

Zhou W, Dai J, Attanasio M, Hildebrandt F. Nephrocystin-3 is required for ciliary function in zebrafish embryos. *Am J Physiol Renal Physiol*. 2010;299:F55-62.